

Engineering, Yeungnam University,
Gyeonsang, Republic of Korea

#Equal contribution

*For correspondence:

jitendrakumarsinha@gmail.com

Received October 24, 2024; Accepted January 31, 2025;
Ahead of print April 02, 2025; Published April 29, 2025

References

- Chakrabarti A, Saha I, Kalita M, Bhattacharya D, Haque B, Nagraj A, *et al*. High prevalence of alcohol use disorder & psychiatric comorbidity among coal mine workers: Observations from a cross-sectional study in West Bengal. *Indian J Med Res* 2024; 160 : 95-101.
- Sedgwick P, Greenwood N. Understanding the Hawthorne effect. *BMJ* 2015; 351 : h4672.
- Rothman KJ, Greenland S, Lash TL. Modern Epidemiology. 3rd ed. Philadelphia: Lippincott Williams & Wilkins; 2008.
- Snipes DJ, Benotsch EG. High-risk cocktails and high-risk sex: Examining the relation between alcohol mixed with energy drinks and sexual risk taking in college students. *Addict Behav* 2013; 38 : 1418-23.
- Seedat S, Scott KM, Angermeyer MC, Berglund P, Bromet EJ, Brugha TS, *et al*. Cross-national associations between gender and mental disorders in the World Health Organization World Mental Health Surveys. *Arch Gen Psychiatry* 2009; 66 : 785-95.
- Tourangeau R, Yan T. Sensitive questions in surveys. *Psychol Bull* 2007; 133 : 859-83.
- Del Boca FK, Darkes J. The validity of self-reports of alcohol consumption: State of the science and challenges for research. *Addiction* 2003; 98 : 1-12.
- Maxwell SE, Cole DA. Bias in cross-sectional analyses of longitudinal mediation. *Psychol Methods* 2007; 12 : 23-44.
- Wang X, Cheng Z. Cross-sectional studies: Strengths, weaknesses, and recommendations. *Chest* 2020; 158 : S65-71.
- Osgood JM, Finan PH, Hinman SJ, So CJ, Quartana PJ. Combat exposure, post-traumatic stress symptoms, and health-related behaviors: The role of sleep continuity and duration. *Sleep* 2019; 42 : zsy257.
- Odes R, Chapman S, Harrison R, Ackerman S, Hong O. Frequency of violence towards healthcare workers in the United States' inpatient psychiatric hospitals: A systematic review of literature. *Int J Ment Health Nurs* 2021; 30 : 27-46.
- Bosma H, Marmot MG, Hemingway H, Nicholson AC, Brunner E, Stansfeld SA. Low job control and risk of coronary heart disease in whitehall II (prospective cohort) study. *BMJ* 1997; 314 : 558-65.
- Edwards AL. The social desirability variable in personality assessment and research. New York: Dryden Press; 1957.
- Shah D. Healthy worker effect phenomenon. *Indian J Occup Environ Med* 2009; 13 : 77-9.
- Cooper ML, Russell M, Skinner JB, Frone MR, Mudar P. Stress and alcohol use: Moderating effects of gender, coping, and alcohol expectancies. *J Abnorm Psychol* 1992; 101 : 139-52.
- Crowne DP, Marlowe D. A new scale of social desirability independent of psychopathology. *J Consult Psychol* 1960; 24 : 349-54.
- Zeng Z, Guo Y, Lu L, Han L, Chen W, Ling L. Mental health status and work environment among workers in small- and medium-sized enterprises in Guangdong, China - a cross-sectional survey. *BMC Public Health* 2014; 14 : 1162.
- Yang L, Wang X, Zhu J, Qin Z. Influencing factors, formation mechanism, and pre-control methods of coal miners' unsafe behavior: A systematic literature review. *Front Public Health* 2022; 10 : 792015.
- Beaton DE, Bombardier C, Guillemin F, Ferraz MB. Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine (Phila Pa 1976)* 2000; 25 : 3186-91.
- O'Cathain A, Hoddinott P, Lewin S, Thomas KJ, Young B, Adamson J, *et al*. Maximising the impact of qualitative research in feasibility studies for randomised controlled trials: Guidance for researchers. *Pilot Feasibility Stud* 2015; 1 : 32.

DOI: 10.25259/IJMR_470_2025

Authors' response

Sir,

Thank you for your interest in our article¹ published in the July 2024 issue of the Indian Journal of Medical Research. We appreciate your thoughtful comments regarding the importance as well as suggested limitations in relation to this paper.

The systematic household-based sampling method was randomized to exclude errors of sampling. Although, limitations, as indicated in the Letter to Editor², could be present, the investigators tried to be statistically precise to avoid such recruitment bias. Similarly, we would have liked to expand our sample size by including more sampling areas. However, keeping the statistical assumptions intact, recruitment was done only from two mining sites, which are representative of the whole population. The focus of the study was exclusively on workers engaged in mining, not on administrative and support staff who do not need to carry out mining work. Therefore, we had to exclude women and other staff as they were not engaged in mining. However, we appreciate the comment that problem areas regarding other mining staff need to be investigated.

Research staff were adequately trained to probe the identification of falsification of information regarding alcohol and other substance use by participants. ASSIST is an internationally validated screening tool for patterns of substance use and provides an accurate measure of patterns of substance use across gender, age and cultures³. We discussed about including biological measures for alcohol and other substance use, but the reported sensitivity and specificity of ASSIST were adequate for identification of the problem at primary care level and we, therefore, decided against using biological measures. The objective of this study was to identify the magnitude of the mental health problem among coal mining workers. Therefore, a cross-sectional design was chosen. The suggestion of a longitudinal study to identify causality is well accepted and shall help in further extrapolation of this study. Anxiety, depression and alcohol use disorders are the three commonest mental health problems, and all can initially be identified and managed at the primary care level. Therefore, we chose the three main problem areas as this population are underserved and have little access to mental healthcare. As a consequence, we did not aim to include other, less common, mental health issues. However, we fully agree that non-inclusion of workplace conditions as predictors of mental health outcomes is a limitation of the study.

Regarding research ethics and informed consent, our research group have a track record of working with mental health issues among different age groups in lower socioeconomic conditions. Therefore, relevant ethical issues of justice and beneficence were duly addressed. Participants identified with 'high risk' alcohol and/or tobacco use and significant anxiety and/or depression were offered coordinated visits to the nearest secondary care treatment facility with mental health services. However, this component was not included in the manuscript, as their visits and outcome could not be monitored.

In conclusion, we appreciate the insights on the limitations of our study. Addressing these issues in future research would strengthen the conclusions and

further enrich our understanding of common mental health issues including alcohol use disorder among underserved populations of India including their causality, predictors and treatment approaches at the primary care level, thereby, further strengthening mental health services at the community level.

Financial support & sponsorship: None.

Conflicts of Interest: None.

Use of Artificial Intelligence (AI)-Assisted Technology for manuscript preparation: The authors confirm that there was no use of AI-assisted technology for assisting in the writing of the manuscript and no images were manipulated using AI.

Amit Chakrabarti^{1,*}, Indranil Saha¹, Manoj Kalita¹, Debangana Bhattacharya¹, Bidisha Haque¹, Alisha Nagraj¹, Sovonlal Mukherjee¹ & Vivek Benegal²

¹ICMR-Centre for Ageing & Mental Health (ICMR-CAM), Kolkata & ²National Institute of Mental Health & Neurosciences (NIMHANS), Bengaluru, India

**For correspondence:*
amit.chakrabarti@gov.in

Received February 20, 2025; Accepted February 20, 2025;
Ahead of print April 02, 2024; Published April 29, 2025

Reference

1. Chakrabarti A, Saha I, Kalita M, Bhattacharya D, Haque B, Nagraj A, *et al.* High prevalence of alcohol use disorder & psychiatric comorbidity among coal mine workers: Observations from a cross-sectional study in West Bengal. *Indian J Med Res* 2024; 160 : 95-101.
2. Ghosh S, Narayanan KB, Singh KK, Sinha JK. Prevalence of alcohol and tobacco use and mental health correlation among male coal miners in West bengal. *Indian J Med Res* 2025; 161 : XX-XX.
3. Humeniuk R, Henry-Edwards S, Ali R., Poznyak V, Monteiro MG. The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): manual for use in primary care. *World Health Organization Geneva*. 2010.